

Code No. 10071

**Anti-Rat
GRO/CINC-1 (6G2) (C) Mouse IgM MoAb**Volume : 500 µg

- Introduction** : Cytokine-induced neutrophil chemo attractant 1(CINC-1) was originally purified from media conditioned by IL-1 β stimulated rat kidney epithelioid cells (NRK-52E). Amino acid sequence that encodes for rat CINC-1 was identified in 1989 by Watanabe's group at Toyama Medical and Pharmaceutical University. CINC-1 is a member of the alpha (CXC) subfamily of chemokines. Three additional rat CXC chemokines (CINC-2 α , CINC-2 β , CINC-3/MIP-2) have been identified. The protein sequence of CINC-1 is 63 - 67% identical to that of CINC-2 α , CINC-2 β , CINC-3/MIP-2. In addition, GRO α , GRO β and GRO γ is sharing 68%, 71% and 69%, identity with CINC-1. This has been suggested that CINCs are the rat counterpart of human GROs.
- Antigen** : Synthetic peptide of the part of the C-terminal of Rat GRO/CINC-1 (CLDPEAPMVQKIVQKMLKGVPK)
- Source** : Mouse-Mouse hybridoma (X63 - Ag 8.653 \times BALB/c mouse spleen cells, [ascites](#))
- Clone** : 6G2 **Subclass** : IgM
- Purification** : Affinity purified by anti-Mouse IgM
- Form** : Lyophilized product in PBS
- How to use** : 1.0 mL deionized water will be added to the product, then its concentration comes to 500 µg/mL
- Stability** : Lyophilized product, 5 years at 2 - 8 °C
: Solution, 2 years at -20 °C
- Application** : This antibody can be used for western blotting in concentration of 5 - 10 µg/mL
- Specificity** : Not react with N-terminal of Rat GRO/CINC-1, Rat GRO/CINC-2 alpha, Rat GRO/CINC-2 beta, Rat GRO/CINC-3, Mouse MIP-2, Human GRO alpha or Human IL-8.
- Reference** : Yagihashi A. et al., Prevention of small intestinal ischemiareperfusion injury in rat by anti-cytokine-induced neutrophil chemoattractant monoclonal antibody. J. of Surg. Res., 1998: 78 (2), 92-96.

For Non-Clinical Research Use Only