

Code No. 10913

**Anti-  
CD134/OX40 (B#11) Mouse IgG MoAb**

Volume : 100 µg

---

**Introduction** : CD134/OX40 is a receptor of OX40 ligand (OX40L) transactivated by Tax of HTLV-1. OX40 is expressed in activated CD4<sup>+</sup> and CD8<sup>+</sup> T cells. OX40 on the activated T cells have a role in proliferation of T cells and production of cytokines by binding with OX40L expressed on antigen presenting cells like dendritic cells or activated B cells. In addition, it also be involved in activated T cell invasion to inflamed area through the binding with OX40L.

**Antigen** : Recombinant soluble OX40

**Source** : Mouse-Mouse hybridoma  
(SP2/0 × BALB/c mouse spleen cells)

**Clone** : B#11

**Subclass** : IgG<sub>1</sub>

**Purification** : Gel filtration chromatography following ammonium sulfate precipitation

**Form** : Lyophilized product from PBS containing 1 % BSA and 0.05 % NaN<sub>3</sub>

**How to use** : 0.2 mL deionized water will be added to the product, then its concentration comes to 500 µg/mL

**Stability** : Lyophilized product, 5 years at 2 - 8 °C  
: Solution, 2 years at -20 °C

**Application** : This antibody can be used for western blotting in concentration of 10 µg/mL.  
: This antibody can be used for immuno-precipitation at 10 µg /test  
: This antibody can be used for flow cytometry analysis at 1 µg (per 1×10<sup>5</sup> cells)  
: This antibody can be used for blocking in concentration of 20 µg/mL  
: This antibody can be used for ELISA (as a detector antibody) in concentration of 5 µg/mL

**Reference** : 1. Kondo K, Okuma K, Tanaka R, Matsuzaki G, Ansari AA, Tanaka Y. Rapid induction of OX40 ligand on primary T cells activated under DNA-damaging conditions. Hum Immunol. 2008 Sep;69(9):533-42.  
2. Kondo K, Okuma K, Tanaka R, Zhang LF, Kodama A, Takahashi Y, Yamamoto N, Ansari AA, Tanaka Y. Requirements for the functional expression of OX40 ligand on human activated CD4<sup>+</sup> and CD8<sup>+</sup> T cells. Hum Immunol. 2007 Jul;68(7):563-71.

---

*For research use only, not for use in diagnostic procedures.*

Code No. 10913

**Anti-  
CD134/OX40 (B#11) Mouse IgG MoAb**

Volume : 500 µg

---

**Introduction** : CD134/OX40 is a receptor of OX40 ligand (OX40L) transactivated by Tax of HTLV-1. OX40 is expressed in activated CD4<sup>+</sup> and CD8<sup>+</sup> T cells. OX40 on the activated T cells have a role in proliferation of T cells and production of cytokines by binding with OX40L expressed on antigen presenting cells like dendritic cells or activated B cells. In addition, it also be involved in activated T cell invasion to inflamed area through the binding with OX40L.

**Antigen** : Recombinant soluble OX40

**Source** : Mouse-Mouse hybridoma  
(SP2/0 × BALB/c mouse spleen cells)

**Clone** : B#11

**Subclass** : IgG<sub>1</sub>

**Purification** : Gel filtration chromatography following ammonium sulfate precipitation

**Form** : Lyophilized product from PBS containing 1 % BSA and 0.05 % NaN<sub>3</sub>

**How to use** : 1.0 mL deionized water will be added to the product, then its concentration comes to 500 µg/mL

**Stability** : Lyophilized product, 5 years at 2 - 8 °C  
: Solution, 2 years at -20 °C

**Application** : This antibody can be used for western blotting in concentration of 10 µg/mL.  
: This antibody can be used for immuno-precipitation at 10 µg /test  
: This antibody can be used for flow cytometry analysis at 1 µg (per 1×10<sup>5</sup> cells)  
: This antibody can be used for blocking in concentration of 20 µg/mL  
: This antibody can be used for ELISA (as a detector antibody) in concentration of 5 µg/mL

**Reference** : 1. Kondo K, Okuma K, Tanaka R, Matsuzaki G, Ansari AA, Tanaka Y. Rapid induction of OX40 ligand on primary T cells activated under DNA-damaging conditions. Hum Immunol. 2008 Sep;69(9) : 533-42.  
2. Kondo K, Okuma K, Tanaka R, Zhang LF, Kodama A, Takahashi Y, Yamamoto N, Ansari AA, Tanaka Y. Requirements for the functional expression of OX40 ligand on human activated CD4<sup>+</sup> and CD8<sup>+</sup> T cells. Hum Immunol. 2007 Jul;68(7) : 563-71.

---

*For research use only, not for use in diagnostic procedures.*

---

*For research use only, not for use in diagnostic procedures.*