

Code No. 18291

Anti-Rat GRO/CINC-3 Rabbit IgG

Volume : 500 μg

Introduction: Growth Related Oncogene (GRO) /Cytokine-induced neutrophil chemo attractant 1

(CINC-1) was originally purified from media conditioned by IL-1βstimulated rat kidney epithelioid cells (NRK-52E). Amino acid sequence that encodes for rat CINC-1 was identified in 1989 by Watanabe's group at Toyama Medical and Pharmaceutical University. CINC-1 is a member of the alpha (CXC) subfamily of chemokines. Three additional rat CXC chemokines (CINC-2α, CINC-2β, CINC-3/MIP-2) have been identified. The protein sequence of CINC-1 is 63 - 67% identical to that of CINC-2α, CINC-2β, CINC-3/MIP-2. In addition, GROα, GROβ and GROγ is sharing 68%, 71% and 69%, identity with CINC-1. This has been suggested that CINCs are the rat

counterpart of human GROs.

Antigen : Recombinant Rat GRO/CINC-3

Purification: Purified with Protein A

Form : Lyophilized product in PBS

How to use: 1.0 mL deionized water will be added to the product (the conc. comes up 500 μg /mL)

Stability: Lyophilized product, 5 years at 2 – 8 °C

: Solution, 2 years at -20 °C

Application: This antibody can be used for western blotting in concentration of 2 - 5 µg /mL.

Specificity: Cross-reacts with mouse MIP-2

Not cross-react with rat GRO/CINC-1, -2α , -2β