


製品番号	製品名			WB	IP	IHC	容量	価格
10343	Anti-	mTOR (N5D11)	Mouse IgG MoAb	約1 µg/mL	3 ~ 5 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28011	Anti-Human	Raptor (R984)	Rabbit IgG Affinity Purify	0.5 ~ 5 µg/mL	1 ~ 5 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28013	Anti-Human	mLST8 (L17)	Rabbit IgG Affinity Purify	0.5 ~ 5 µg/mL	1 ~ 5 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28015	Anti-Human	Rheb (R165)	Rabbit IgG Affinity Purify	0.5 ~ 5 µg/mL	1 ~ 5 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28033	Anti-Human	PRAS40 (P238)	Rabbit IgG Affinity Purify	0.5 ~ 3 µg/mL	1 ~ 3 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28035	Anti-Human	PRAS40 (S183 Phosphorylated)	Rabbit IgG Affinity Purify	1 ~ 3 µg/mL	-	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28037	Anti-Human	Raptor (R1)	Rabbit IgG Affinity Purify	1 ~ 3 µg/mL	0.25 ~ 3 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28039	Anti-Human	Rictor (R1291)	Rabbit IgG Affinity Purify	1 ~ 3 µg/mL	1 ~ 3 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28017	Anti-Human	TSC1/Hamartin (T1149)	Rabbit IgG Affinity Purify	1 ~ 5 µg/mL	-	1 ~ 5 µg/mL	100 µg	¥ 58,000
							10 µg	¥ 15,000
28063	Anti-Human	GBF1 (1844)	Rabbit IgG Affinity Purify	1 ~ 5 µg/mL	1 ~ 5 µg/test	-	100 µg	¥ 58,000
							10 µg	¥ 15,000
28065	Anti-Human	GBF1 (T1337 Phosphorylated)	Rabbit IgG Affinity Purify	1 ~ 5 µg/mL	-	-	100 µg	¥ 58,000
							10 µg	¥ 15,000

参考文献 :

- Miyamoto T, Oshiro N, Yoshino K, Nakashima A, Eguchi S, Takahashi M, Ono Y, Kikkawa U, Yonezawa K. AMP-activated Protein Kinase Phosphorylates Golgi-specific Brefeldin A Resistance Factor 1 at Thr1337 to Induce Disassembly of Golgi Apparatus. *J Biol Chem.* 2008 Feb 15;283(7):4430-8.
- Nakashima A, Yoshino K, Miyamoto T, Eguchi S, Oshiro N, Kikkawa U, Yonezawa K. Identification of TBC7 having TBC domain as a novel binding protein to TSC1-TSC2 complex. *Biochem Biophys Res Commun.* 2007 Sep 14;361(1):218-23.
- Yang Q, Guan KL. Expanding mTOR signaling. *Cell Res.* 2007 Aug;17(8):666-81.
- Oshiro N, Takahashi R, Yoshino K, Tanimura K, Nakashima A, Eguchi S, Miyamoto T, Hara K, Takehana K, Avruch J, Kikkawa U, Yonezawa K. The proline-Rich Akt substrate of 40 kDa (PRAS40) is a physiological substrate of mTOR complex 1. *J Biol Chem.* 2007 Jul 13;282(28):20329-39.
- Avruch J, Hara K, Lin Y, Liu M, Long X, Ortiz-Vega S, Yonezawa K. Insulin and amino-acid regulation of mTOR signaling and kinase activity through the Rheb GTPase. *Oncogene.* 2006 Oct 16;25(48):6361-72.
- Takahara T, Hara K, Yonezawa K, Sorimachi H, Maeda T. Nutrient-dependent multimerization of the mammalian target of rapamycin through the N-terminal HEAT repeat region. *J Biol Chem.* 2006 Sep 29;281(39):28605-14.
- Eguchi S, Tokunaga C, Hidayat S, Oshiro N, Yoshino K, Kikkawa U, Yonezawa K. Different roles for the TOS and RAIP motifs of the translational regulator protein 4E-BP1 in the association with raptor and phosphorylation by mTOR in the regulation of cell size. *Genes Cells.* 2006 Jul;11(7):757-66.
- Ojiji G, Hidayat S, Nakashima A, Tokunaga C, Oshiro N, Yoshino K, Yokono K, Kikkawa U, Yonezawa K. Suppression of the mTOR-raptor signaling pathway by the inhibitor of heat shock protein 90 geldanamycin. *J Biochem (Tokyo).* 2006 Jan;139(1):129-35.
- Yonezawa K. Rapamycin. *Nippon Rinsho.* 2005 May;63 Suppl 5:718-22.
- Long X, Lin Y, Ortiz-Vega S, Yonezawa K, Avruch J. Rb binds and regulates the mTOR kinase. *Curr Biol.* 2005 Apr 26;15(8):702-13.
- Nakajyo T, Yamatsuji T, Ban H, Shigemitsu K, Haisa M, Motoki T, Noma K, Nobuhisa T, Matsuoaka J, Gunduz M, Yonezawa K, Tanaka N, Naomoto Y. Glutamine is a key regulator for amino acid-controlled cell growth through the mTOR signaling pathway in rat intestinal epithelial cells. *Biochem Biophys Res Commun.* 2005 Jan 7;326(1):174-80.
- Takei N, Inamura N, Kawamura M, Namba H, Hara K, Yonezawa K, Nawa H. Brain-derived neurotrophic factor induces mammalian target of rapamycin-dependent local activation of translation machinery and protein synthesis in neuronal dendrites. *J Neurosci.* 2004 Nov 3;24(44):9760-9.
- Murakami M, Ichisaka T, Maeda M, Oshiro N, Hara K, Edenhofer F, Kiyama H, Yonezawa K, Yamanaka S. mTOR is essential for growth and proliferation in early mouse embryos and embryonic stem cells. *Mol Cell Biol.* 2004 Aug;24(15):6710-8.
- Oshiro N, Yoshino K, Hidayat S, Tokunaga C, Hara K, Eguchi S, Avruch J, Yonezawa K. Dissociation of raptor from mTOR is a mechanism of rapamycin-induced inhibition of mTOR function. *Genes Cells.* 2004 Apr;9(4):359-66.
- Tokunaga C, Yoshino K, Yonezawa K. mTOR integrates amino acid- and energy-sensing pathways. *Biochem Biophys Res Commun.* 2004 Jan 9;313(2):443-6.
- Yonezawa K, Tokunaga C, Oshiro N, Yoshino K. Raptor, a binding partner of target of rapamycin. *Biochem Biophys Res Commun.* 2004 Jan 9;313(2):437-41.
- Yonezawa K, Yoshino KI, Tokunaga C, Hara K. Kinase activities associated with mTOR. *Curr Top Microbiol Immunol.* 2004;279:271-82.
- Yoshino K, Tokunaga C, Hara K, Yonezawa K. mTOR complex as an integrating controller of cell growth. *Tampakushti Kakusan Koso.* 2003 Aug;48(10):1378-85.
- Gao X, Pan D. TSC1 and TSC2 tumor suppressors antagonize insulin signaling in cell growth. *Genes Dev.* 2001 Jun 1;15(11):1383-92.
- Kobayashi T, Minowa O, Sugitani Y, Takai S, Mitani H, Kobayashi E, Noda T, Hino O. A germ-line Tsc1 mutation causes tumor development and embryonic lethality that are similar, but not identical to, those caused by Tsc2 mutation in mice. *Proc Natl Acad Sci U S A.* 2001 Jul 17;98(15):8762-7.
- Nishiumi T, Hara K, Tsujishita Y, Kaneko K, Shii K, Yonezawa K. Characterization of the phosphoproteins and protein kinase activity in mTOR immunoprecipitates. *Biochem Biophys Res Commun.* 1998 Nov 18;252(2):440-4.
- Plank TL, Yeung RS, Henske EP. Hamartin, the product of the tuberous sclerosis 1 (TSC1) gene, interacts with tuberin and appears to be localized to cytoplasmic vesicles. *Cancer Res.* 1998 Nov 1;58(21):4766-70.
- van Slegtenhorst M, de Hoogt R, Hermans C, Nellist M, Janssen B, Verhoef S, Lindhout D, van den Ouwehand A, Halley D, Young J, Burley M, Jeremiah S, Woodward K, Nahmias J, Fox M, Ekong R, Osborne J, Wolfe J, Povey S, Snell RG, Cheadle JP, Jones AC, Tachataki M, Ravine D, Sampson JR, Reeve MP, Richardson P, Wilmer F, Munro C, Hawkins TL, Sepp T, Ali JB, Ward S, Green AJ, Yates JR, Kwiatkowska J, Henske EP, Short MP, Haines JH, Jozwiak S, Kwiatkowski DJ. Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34. *Science.* 1997 Aug 8;277(5327):805-8.
- European Chromosome 16 Tuberous Sclerosis Consortium. Identification and characterization of the tuberous sclerosis gene on chromosome 16. *Cell.* 1993 Dec 31;75(7):1305-15.

ラパマイシンは免疫抑制効果、抗がん効果、心臓冠状動脈治療における再狭窄防止効果等を有する、マクロライド系有機化合物で、mTOR (Mammalian target of rapamycin) はラパマイシンの細胞内標的として見つかった、約 290kDa のセリン / スレオニンキナーゼです。近年 mTOR は細胞成長における重要な役割と腫瘍形成への関わりで非常に注目されており、種々の mTOR 関連タンパクも同定され、mTOR を取り巻くシグナル制御のネットワークが解明されつつあります。



取扱い販売代理店