

Code No. 18115

**Anti-Human  
SCF (K165) Rabbit IgG Affinity Purify**

Volume : 100 µg

---

**Introduction** : Stem cell factor (SCF) and its receptor KIT play an important role in various biologic phases, such as hematopoiesis, reproduction, and regeneration. Structurally, c-Kit contains five immunoglobulin-like domains extracellularly and a catalytic domain divided into two regions by a 77 amino acid insert intracellularly. Studies in white spotting and steel mice have shown that functional SCF and c-Kit are critical in the survival and development of stem cells involved in hematopoiesis, pigmentation and reproduction. Mutations in c-Kit are associated with a variety of human diseases. Interaction of SCF with c-Kit rapidly induces receptor dimerization and increases in autophosphorylation activity. Downstream of c-Kit, multiple signal transduction components are activated, including phosphatidylinositol-3-kinase, Src family members, the JAK/STAT pathway and the Ras-Raf-MAP kinase cascade.

**Antigen** : Synthetic peptide of a part of Human SCF

**Purification** : Purified with antigen peptide

**Form** : Lyophilized product from 1 % BSA in PBS containing 0.05% NaN<sub>3</sub>

**How to use** : 1.0 mL deionized water will be added to the product (the conc. comes up 100 µg /mL)

**Stability** : Lyophilized product, 5 years at 2 – 8 °C  
: Solution, 2 years at –20 °C

**Application** : This antibody can be used for immunohistochemistry with formalin fixed paraffin embedded tissues by several techniques such as Avidin Biotin Complex (ABC) Method. The optimal concentration is about 1 - 5 µg/mL, however, the concentration should be optimized by each laboratory.  
: This antibody can be used for western blotting in concentration of 2 - 5 µg /mL.