

Code No. 18903

**Anti-Human
GLUT-3 Rabbit IgG Affinity Purify**

Volume : 100 µg

GLUT : Glucose Transporter

Introduction : Mammalian cells transport glucose through a family of membrane proteins known as glucose transporters (GLUTs or SLC2A family). Mammalian tissues are known to express different glucose transporter isoforms (existing at least GLUTs 1-13) in different tissues. The molecular weight are 40-60 kDa, respectively. GLUT-3 has a high affinity with glucose (Km 1 -5 mM) and is the main transporter in brain (neurons).

Antigen : Synthetic peptide of the C terminal part of human GLUT-3 (SGKDGVMEMNSIEPAKETT)

Purification : Purified with antigen peptide

Form : Lyophilized product from PBS containing 1 % BSA and 0.05% NaN₃

How to use : 1.0 mL deionized water will be added to the product (the conc. comes up 100 µg /mL)

Stability : Lyophilized product, 5 years at 2 – 8 °C
: Solution, 2 years at –20 °C

Application : This antibody can be used for immunohistochemistry with formalin fixed paraffin embedded tissues without any pretreatment by several techniques such as Avidin Biotin Complex (ABC) Method. The optimal concentration is 2-5 µg/mL, however, the concentration should be optimized by each laboratory.
: This antibody can be used for western blotting in concentration of 1 - 5 µg /mL.

Specificity : Confirmed by western blotting using transfectant cells.

For research use only, not for use in diagnostic procedures.

Code No. 18903

**Anti-Human
GLUT-3 Rabbit IgG Affinity Purify**

Volume : 10 µg

GLUT : Glucose Transporter

Introduction : Mammalian cells transport glucose through a family of membrane proteins known as glucose transporters (GLUTs or SLC2A family). Mammalian tissues are known to express different glucose transporter isoforms (existing at least GLUTs 1-13) in different tissues. The molecular weight are 40-60 kDa, respectively. GLUT-3 has a high affinity with glucose (Km 1 -5 mM) and is the main transporter in brain (neurons).

Antigen : Synthetic peptide of the C terminal part of human GLUT-3 (SGKDGVMEMNSIEPAKETT)

Purification : Purified with antigen peptide

Form : Lyophilized product from PBS containing 1 % BSA and 0.05% NaN₃

How to use : 0.1 mL deionized water will be added to the product (the conc. comes up 100 µg /mL)

Stability : Lyophilized product, 5 years at 2 – 8 °C
: Solution, 2 years at –20 °C

Application : This antibody can be used for immunohistochemistry with formalin fixed paraffin embedded tissues without any pretreatment by several techniques such as Avidin Biotin Complex (ABC) Method. The optimal concentration is 2-5 µg/mL, however, the concentration should be optimized by each laboratory.
: This antibody can be used for western blotting in concentration of 1 - 5 µg /mL.

Specificity : Confirmed by western blotting using transfectant cells.

For research use only, not for use in diagnostic procedures.