

Code No. 28041

Anti-GZF1 Rabbit IgG Affinity Purify

Volume : 100 µg

Introduction	:	<i>GZF1</i> has been identified as a gene whose expression is induced by a neurotrophic factor, GDNF. It codes a protein which has BTB/POZ domain in its N-terminal region and 10 of zinc finger motifs in its C-terminal area from the center. It has been known by performance analysis that GZF1 is a transcriptional repression factor which sequence-specifically binds to DNA and the binding sequence is present in transcription control region of HOXA10.
Antigen	:	Synthetic peptide of the part of Human GZF1
Purification	:	Purified with antigen peptide
Form	:	Lyophilized product from 1% BSA in PBS containing 0.05% NaN $_3$
How to use	:	1.0 mL deionized water will be added to the product (the conc. comes up 100 μg /mL)
Stability	:	Lyophilized product, 5 years at 2 - 8 °C Solution, 2 years at –20 °C
Application	:	This antibody can be used for western blotting in concentration of about 0.1 μ g /mL. This antibody can be used for immuno-precipitation in concentration about 1 μ g /mL.
Specificity	:	React with human and mouse GZF1
Reference	:	 Fukuda N, Ichihara M, Morinaga T, Kawai K, Hayashi H, Murakumo Y, Matsuo S, Takahashi M. Identification of a novel glial cell line-derived neurotrophic factor-inducible gene required for renal branching morphogenesis. J Biol Chem. 2003 Dec 12;278(50):50386-92. Morinaga T, Enomoto A, Shimono Y, Hirose F, Fukuda N, Dambara A, Jijiwa M, Kawai K, Hashimoto K, Ichihara M, Asai N, Murakumo Y, Matsuo S, Takahashi M. GDNF-inducible zinc finger protein 1 is a sequence-specific transcriptional repressor that binds to the HOXA10 gene regulatory region. Nucleic Acids Res. 2005 Jul 26;33(13):4191-201.

For Non-Clinical Research Use Only